THE EXISTENCE OF MULTIPLE SOLUTIONS FOR BOUNDARY VALUE PROBLEM WITH ONE DIMENSIONAL P-LAPLACIAN

BORIS RUDOLF

ABSTRACT. The boundary value problem for a differential equation with one dimensional p-Laplacian is studied. The technique of lower and upper solutions is used. The existence of a solution for well ordered and unordered case as well as the existence of multiple solutions are proved. The growth condition is assumed only on a part of the nonlinearity.

The paper deals with the boundary value problem

$$(\varphi_p(x'))' = F(t, x, \varphi_p(x')), \quad (1)$$

$$x(b) = \int_0^b x(s) dg(s) - k\varphi_p(x'(b)), \quad x'(0) = 0. \quad (2)$$

The left hand side of (1) is one dimensional p-Laplacian, $\varphi_p(s) = |s|^{p-1} \operatorname{sgn}(s)$, the right hand side is the sum of two continuous functions

$$F(t, x, y) = f_1(t, x, y) + f_2(t, x, y)$$

with different properties. The second boundary condition is nonlocal, $g(s)$ is a nondecreasing function of bounded variation, $G(s) = \operatorname{var}_{[0,s]} g(\tau)$, $G(b) < 1$ and $k \geq 0$.

The nonlocal boundary condition covers certain types of linear two point, multipoint and integral boundary conditions.

The main goal is to prove existence of multiple classical solutions of (1), (2), i.e., solutions from

$$D = \{x \in C^1(I), \varphi_p(x') \in C^1(I)\}, \quad I = [0, b]$$

under suitable conditions.

© 2013 Mathematical Institute, Slovak Academy of Sciences.

2010 Mathematics Subject Classification: 34B10, 34B15.

Keywords: p-Laplacian, nonlocal boundary condition, lower and upper solution.

Supported by grant 1/0021/10 of the Scientific Grant Agency VEGA of Slovak Republic.

We use the method of lower and upper solutions. Results of this type for the classical second order BVP are well known, see [4], [7], existence results for periodic BVP with p-Laplacian and growth condition on nonlinearity are in [5], and existence result for (1), (2) in well ordered case and restricted growth of nonlinearity is given in [8].

The paper is based on ideas of [7] applied on the problem (1), (2). By a priori estimates of the norm of solutions or its derivatives we use method of Korman [3], based on Bihari inequality [1].

We define a lower and upper solution of (1), (2) by the obvious way. Set $I^0 = I \setminus \{t_i; 0 < t_i < b, i = 1 \dots n\}$.

DEFINITION 1 ([8]). A function $\alpha \in C(I) \cup C^1(I^0)$, with $\varphi(\alpha') \in C^1(I^0)$ is called a lower solution of (1), (2) if

$$\begin{aligned} \lim_{t \rightarrow t_i -} \alpha'(t) &\leq \lim_{t \rightarrow t_i +} \alpha'(t) && \text{for } i = 1, \dots, n, \\ \left(\varphi_p(\alpha'(t)) \right)' &\geq f\left(t, \alpha(t), \varphi_p(\alpha'(t))\right) && \text{for } t \in I^0, \\ \alpha(b) &\leq \int_0^b \alpha(s) dg(s) - k\varphi_p(\alpha'(b)), \quad \alpha'(0) \geq 0. \end{aligned}$$

Similarly a function $\beta \in C(I) \cup C^1(I^0)$, with $\varphi(\beta') \in C^1(I^0)$ is called an upper solution of (1), (2) if

$$\begin{aligned} \lim_{t \rightarrow t_i -} \beta'(t) &\geq \lim_{t \rightarrow t_i +} \beta'(t) && \text{for } i = 1, \dots, n, \\ \left(\varphi_p(\beta'(t)) \right)' &\leq f\left(t, \beta(t), \varphi_p(\beta'(t))\right) && \text{for } t \in I^0, \\ \beta(b) &\geq \int_0^b \beta(s) dg(s) - k\varphi_p(\beta'(b)), \quad \beta'(0) \leq 0. \end{aligned}$$

In the case of strict inequalities for the equation on I^0 and for the second boundary condition we say that lower and upper solutions are strict.

LEMMA 2 ([8]). *Let α, β be a strict lower and upper solution and x be a solution of the problem (1), (2). Then $\alpha(t) \leq x(t)$ for each $t \in I$ implies $\alpha(t) < x(t)$ for each $t \in I$ and $\beta(t) \geq x(t)$ for each $t \in I$ implies $\beta(t) > x(t)$ for each $t \in I$.*

In our first existence result as well as in all theorems below we assume that the summand f_1 satisfies a Nagumo–Bernstein type of growth condition [2], [5] and f_2 instead of this satisfies one of two possible types of sign condition.

In the rest of the paper constant $q > 0$ is such that $\frac{1}{p} + \frac{1}{q} = 1$. Then the function φ_q is the inverse of φ_p .

THEOREM 3. *Let*

(A1) $\exists r > 0$ such that $\forall t \in I$, $f_1(t, r, 0) > 0$, $f_1(t, -r, 0) < 0$ on I ,

(B1) $\exists c > 0$ such that $\forall t \in I$, $|x| < r$, $y \in R$,

$$|f_1(t, x, y)| \leq c(1 + |y|^q),$$

(C1) $\forall t \in I$, $x, y \in R$,

$$x \cdot f_2(t, x, y) \geq 0.$$

Then the problem (1), (2) possesses a solution x such that $|x(t)| < r$.

PROOF. Set $X = C^1([0, b])$ and define an operator $T: X \rightarrow X$ by

$$Tx(t) = \frac{1}{G(b) - 1} \left\{ \int_0^b G(s) \varphi_q(\tilde{F}_x(s)) ds + k(\tilde{F}_x(b)) \right\} - \int_t^b \varphi_q(\tilde{F}_x(s)) ds, \quad (3)$$

where

$$\tilde{F}_x(s) = \int_0^s F(\tau, x(\tau), \varphi_p(x'(\tau))) d\tau.$$

Then

$$Tx(t) \in D = \{x \in C^1(I), \varphi_p(x') \in C^1(I)\}, \quad \text{and} \quad (Tx)'(0) = 0.$$

The function satisfies boundary conditions (2), see [8], and fixed point of completely continuous operator T is a solution of (1), (2).

We consider the perturbed boundary value problems

$$(\varphi_p(x'))' = \lambda F(t, x, \varphi_p(x')) + (1 - \lambda)x(t), \quad (4)$$

$$x(b) = \int_0^b x(s) dg(s) - k\varphi_p(x'(b)), \quad x'(0) = 0 \quad (5)$$

with $\lambda \in [0, 1]$.

Clearly, $-r$ and r are strict lower and upper solutions of (4), (5).

Convex combination $f_{1\lambda} = \lambda f_1 + (1 - \lambda)x$ satisfies the same growth condition (A1) with the constant $c_1 = c + r$ and $f_{2\lambda} = \lambda f_2$ satisfies the sign condition (C1) independently on λ .

Let $x(t)$ be a solution of (4), (5) with $|x(t)| < r$ on I . We estimate $|x'(t)|$.

Suppose $x(t_0) = \max_{[t_0, t_1]}$, $x(t) > 0$ and $x'(t) < 0$ on (t_0, t_1) . Then $x(t)$ is invertible on $[t_0, t_1]$, its inverse is $t(x)$. Denote $p(x) = \varphi_p(x'(t(x)))$.

We rewrite the equation (4) in

$$p'(x)\varphi_q(p(x)) = f_{1\lambda}(t, x, p(x)) + f_{2\lambda}(t, x, p(x))$$

and estimate

$$p'(x)\varphi_q(p(x)) \geq -c_1(1 + |p(x)|^q).$$

Substitution $z(x) = |p(x)|^q$ gives

$$z'(x) \geq -qc_1(1 + z(x))$$

and after the change of variable $v = x_0 - x$ we denote $\tilde{z}(v) = z(x) + 1$ and obtain

$$\tilde{z}'(v) \leq qc_1\tilde{z}(v), \quad \tilde{z}(0) = 1.$$

Then

$$\tilde{z}(v) \leq e^{qc_1v} \leq e^{qc_1r} \quad \text{and} \quad |p(x)| \leq e^{c_1r} = c_2.$$

That means, $-\varphi_q(c_2) \leq x'(t) \leq 0$ on $[t_0, t_1]$. Boundedness of derivative in case $x > 0$ and $x' > 0$ and in cases $x < 0$ is proved similarly. Then boundedness of a solution x , $x(t) < r$ implies the existence of a constant ρ , $x'(t) < \rho$ on I .

We set

$$\Omega_{r,\rho} = \{x \in X; |x| < r, |x'| < \rho\}.$$

The associated homotopy operator

$$H(x, \lambda) = \frac{1}{G(b) - 1} \left\{ \int_0^b G(s) \varphi_q(\tilde{F}_{x,\lambda}(s)) ds + k(\tilde{F}_{x,\lambda}(b)) \right\} - \int_t^b \varphi_q(\tilde{F}_{x,\lambda}(s)) ds,$$

with $\tilde{F}_{x,\lambda}(s) = \int_0^s \lambda f_1(\tau, x(\tau), \varphi_p(x'(\tau))) + f_2(\tau, x(\tau), \varphi_p(x'(\tau))) + (1-\lambda)x(\tau) d\tau$ possesses no fixed point on the boundary of $\Omega_{r,\rho}$. Then the Leray-Schauder degree of $H(., \lambda)$ is well defined and independent on λ .

$H(x, 0)$ is an odd operator and Borsuk's theorem implies [2]

$$d(I - T, \Omega, 0) = d(I - H(x, 0), \Omega, 0) = 1 \pmod{2}$$

which implies the existence of a fixed point $x \in \Omega$ of T . □

By the same method can be proved the following existence result with another sign condition.

THEOREM 4. *Let*

(A1) $\exists r > 0$ such that $\forall t \in I$, $f_1(t, r, 0) > 0$, $f_1(t, -r, 0) < 0$ on I ,

(B1) $\exists c > 0$ such that $\forall t \in I$, $|x| < r$, $y \in R$,

$$|f_1(t, x, y)| \leq c(1 + |y|^q),$$

(C2) $\forall t \in I$, $x, y \in R$,

$$y \cdot f_2(t, x, y) \leq 0.$$

Then the problem (1), (2) possesses a solution x such that $|x(t)| < r$.

In the case of well ordered nonconstant lower and upper solutions the following theorem holds.

THEOREM 5. *Let*

(A2) $\alpha(t) \leq \beta(t)$ *be a lower and upper solution of (1), (2),*

(B2) $\exists c(r) : R^+ \rightarrow R^+$ *such that* $\forall t \in I, |x| < r, y \in R,$

$$|f_1(t, x, y)| \leq c(r)(1 + |y|^q),$$

(C1) or (C2).

Then the problem (1), (2) possesses a solution x such that $\alpha(t) \leq x(t) \leq \beta(t)$.

PROOF. We prove the case when (C1) holds. Set $r = \max\{|\alpha|, |\beta|\} + 1$, and choose $M > \max\{|f(t, x, 0)|; t \in I, \alpha(t) \leq x \leq \beta(t)\}$.

Define

$$f_1^*(t, x, y) = \begin{cases} f_1(t, \beta(t), y) + M(r - \beta(t)), & x > r, \\ f_1(t, \beta(t), y) + M(x - \beta(t)), & \beta(t) < x \leq r, \\ f_1(t, x, y), & \alpha(t) \leq x \leq \beta(t), \\ f_1(t, \alpha(t), y) - M(\alpha(t) - x), & -r \leq x < \alpha(t), \\ f_1(t, \alpha(t), y) - M(\alpha(t) + r), & x < -r, \end{cases}$$

and

$$f_2^*(t, x, y) = \begin{cases} f_2(t, \beta(t), y), & 0 \leq \beta(t) < x, \\ f_2(t, \beta(t), y) \frac{x}{\beta(t)}, & \beta(t) < x < 0, \\ 0, & \beta(t) < 0 \leq x, \\ f_2(t, x, y), & \alpha(t) \leq x \leq \beta(t), \\ 0, & x \leq \alpha(t) < 0, \\ f_2(t, \alpha(t), y) \frac{x}{\alpha(t)}, & 0 < x < \alpha(t), \\ f_2(t, \alpha(t), y), & x < \alpha(t) \leq 0. \end{cases}$$

Now the boundary value problem

$$(\varphi_p(x'))' = f_1^*(t, x, \varphi_p(x')) + f_2^*(t, x, \varphi_p(x')), \quad (6)$$

$$x(b) = \int_0^b x(s) dg(s) - k\varphi_p(x'(b)), \quad x'(0) = 0 \quad (7)$$

satisfies conditions (A1), (B1), and (C1). That means, perturbed BVP is solvable and

$$d(I - T^*, \Omega_{r,p}, 0) = 1 \pmod{2}.$$

Moreover, for each $\epsilon > 0$ the function $\alpha(t) - \epsilon$ is a strict lower solution and $\beta(t) + \epsilon$ is a strict upper solution of (6), (7).

Lemma 2 implies that $\alpha(t) \leq x(t) \leq \beta(t)$. That means, $f_i^*(t, x, \varphi_p(x')) = f_i(t, x, \varphi_p(x'))$ and $x(t)$ is also a solution of (1), (2). $\square$

EXAMPLE 6. We consider the boundary value problem

$$\begin{aligned} (\varphi_p(x'))' + k(t, x)\varphi_m(x') + f(t, x) &= h(t), \\ x(1) &= \int_0^1 x(s) dg(s) - k\varphi_p(x'(1)), \quad x'(0) = 0, \end{aligned} \quad (8)$$

and assume that

$$k(t, x) \geq 0, \quad m > 1, \quad \text{and} \quad \lim_{x \rightarrow \infty} f(t, x) = -\infty, \quad \lim_{x \rightarrow -\infty} f(t, x) = \infty.$$

Then BVP (8) possesses a solution for each $h(t) \in C(I)$.

EXAMPLE 7. Consider the boundary value problem

$$\begin{aligned} (\varphi_p(x'))' + k(t, x)\varphi_r(x') &= f(t, x, x'), \\ x(1) &= \int_0^1 x(s) dg(s) - k\varphi_p(x'(1)), \quad x'(0) = 0 \end{aligned} \quad (9)$$

with the bounded nonlinearity $|f(t, x, y)| \leq M$, and $k(t, x) \geq 0$. Then BVP (9) possesses a solution $x(t)$,

$$|x(t)| \leq \frac{1}{1 - G(1)} (B_0 + kMq^{1-p}) + \beta_0(t),$$

where

$$\beta_0(t) = Mq^{1-p}(1 - t^q) \quad \text{and} \quad B_0 = \int_0^1 \beta(s) dg(s).$$

The case of unordered lower and upper solutions is more complicated, requires stronger growth conditions on f_1 . The existence result presented below gives no a priori bound of a solution, instead of this we have only a partial information about its localization.

THEOREM 8. *Let*

(A3) $\alpha \not\leq \beta$ *be a strict lower and upper solution of (1), (2),*

(B3) $\exists 0 < A, B < q - 1, c > 0$ *such that $\forall t \in I, x, y \in R,$*

$$|f_1(t, x, y)| \leq c(1 + |x|^A + |y|^B),$$

(C1) or (C2).

Then the problem (1), (2) possesses a solution x such that $\beta \not\leq x \not\leq \alpha$.

Proof. We set $r_0 = \max(|\alpha|, |\beta|)$, a number $r > r_0$, we specify later, and $M_0 = c(1 + r^A)$. We define a perturbation

$$f_1^*(t, x, y) = \begin{cases} f_1(t, r, y) + M_0, & x > r + 1, \\ f_1(t, r, y) + M_0(x - r), & r < x \leq r + 1, \\ f_1(t, x, y), & -r \leq x \leq r, \\ f_1(t, r, y) + M_0(x + r), & -r - 1 \leq x < -r, \\ f_1(t, r, y) - M_0, & x < -r - 1. \end{cases}$$

Clearly, $-r - 1$, α , β , $r + 1$ are strict lower and upper solutions of the perturbed BVP

$$(\varphi_p(x'))' = f_1^*(t, x, \varphi_p(x')) + f_2(t, x, \varphi_p(x')), \quad (10)$$

$$x(b) = \int_0^b x(s) dg(s) - k\varphi_p(x'(b)), \quad x'(0) = 0. \quad (11)$$

Moreover, f_1^* satisfies the growth property (B1) and (B2) for chosen r

$$|f_1^*(t, x, y)| \leq c(1 + |r|^A + |y|^B) + M_0 < \bar{c}(1 + |y|^q).$$

The existence theorems above and their proofs imply (T^* is given by (3) using f_1^* instead of f)

$$d(I - T^*, \Omega_{r+1, \rho}, 0) = d(I - T^*, \Omega_{\alpha, r+1, \rho}, 0) = d(I - T^*, \Omega_{-r-1, \beta, \rho}, 0) = 1 \pmod{2}.$$

Then

$$d(I - T^*, \Omega, 0) = 1 \pmod{2} \quad \text{for } \Omega = \Omega_{r+1, \rho} \setminus \overline{\Omega_{\alpha, r+1, \rho} \cup \Omega_{-r-1, \beta, \rho}}.$$

We estimate the norm of a solution x using a method of Korman [3], [7]. We consider the case (C1) only.

Suppose that $\max |x(t)| = x(t_0) > r_0$ and set $M = x(t_0) - r_0 > 0$. The second boundary condition implies $t_0 \neq b$. As $x \in \Omega$, $\exists t_\alpha$ such that $x(t_\alpha) \leq \alpha(t_\alpha) \leq r_0$. Then $\exists t_1$ such that $x(t_1) = r_0$.

We suppose $t_0 < t_1$, the opposite case is treated similarly.

As in the proof of Theorem 3, we denote $p(x) = \varphi_p(x'(t(x)))$, $z(x) = |p(x)|^q$ and $v = M + r_0 - x$, to obtain

$$\tilde{z}'(v) \leq qc \left(1 + (r + 1)^A + \tilde{z}(v)^{\frac{B}{q}} \right), \quad \tilde{z}(0) = 0.$$

The Bihari lemma [1] implies

$$\tilde{z}(v) \leq \left[\left(cqM(1 + (r + 1)^A) \right)^{1 - \frac{B}{q}} + (q - B)cM \right]^{\frac{1}{1 - \frac{B}{q}}}.$$

Then

$$|p(x)| \leq \left[\left(cqM(1 + (r+1)^A) \right)^{1-\frac{B}{q}} + (q-B)cM \right]^{\frac{1}{q-B}}$$

and

$$|p(x)| \leq d \left(\left(cqM(1 + (r+1)^A) \right)^{\frac{1}{q}} + ((q-B)cM)^{\frac{1}{q-B}} \right), \quad d = 2^{\frac{1}{q-B}}.$$

As $M \leq (b-a) \max_{v \in [0, M]} |p(v)|$, we have

$$M \leq d(b-a) \left(\left(cqM(1 + (r+1)^A) \right)^{\frac{1}{q}} + ((q-B)cM)^{\frac{1}{q-B}} \right).$$

Then

$$M^{\frac{q-1}{q}} \leq c_1 r^{\frac{A}{q}} + c_2 M^{\frac{b}{(q-B)q}},$$

where positive constants c_i are independent on r .

Either

$$r_0 + M < r \quad \text{or} \quad r_0 + M \geq r.$$

In the latter case

$$M^{\frac{q-1}{q}} \leq c_1 (r_0 + M)^{\frac{A}{q}} + c_2 M^{\frac{b}{(q-B)q}}.$$

Then $\exists M_1$ independent of r such that $r_0 + M < r$.

We proceed similarly also in the case $\max |x(t)| = -x(t_0) > r_0$. That means, $|x(t)| < r$ and then x is a solution of the original problem. $\square$

We prove the existence of at least two solutions under assumption that the boundary value problem has either two strict lower solutions and one upper one or vice versa.

LEMMA 9. *Let α be a strict lower solution of (1), (2).*

Suppose (C1) holds and sets

$$f_{1\alpha}(t, x, y) = \begin{cases} f_1(t, x, y) & \text{for } t \in I, x \geq \alpha(t), y \in R, \\ f_1(t, \alpha(t), y) & \text{for } t \in I, x < \alpha(t), y \in R, \end{cases} \quad (12)$$

and

$$f_{2\alpha}(t, x, y) = \begin{cases} f_2(t, x, y) & \text{for } t \in I, x \geq \alpha(t), y \in R, \\ x f_2(t, \alpha(t), y) & \text{for } t \in I, 0 \leq x < \alpha(t), y \in R, \\ 0 & \text{for } t \in I, x < 0 \leq \alpha(t), y \in R, \\ f_2(t, \alpha(t), y) & \text{for } t \in I, x < \alpha(t) < 0, y \in R. \end{cases} \quad (13)$$

Then each solution $x(t)$ of the problem

$$\begin{aligned} (\varphi_p(x'))' &= F_\alpha(t, x, \varphi_p(x')), \\ x(b) &= \int_0^b x(s) dg(s) - k\varphi_p(x'(b)), \quad x'(0) = 0 \end{aligned} \quad (14)$$

is a solution of (1), (2).

Proof. The proof is based on the fact that in both cases the function $\alpha(t) - m$ is a strict lower solution of (14) for each $m > 0$. Then Lemma 2 implies our assertion. $\square$

THEOREM 10. *Let*

(A4) $\alpha \leq \beta$, $\alpha \leq \alpha_1$, $\alpha_1 \not\leq \beta$, α, α_1 be a strict lower and β a strict upper solution of (1), (2),

(B4) $\exists 0 < A, B < q - 1$, $c > 0$ such that $\forall t \in I$, $x, y \in R$,

$$f_1(t, x, y) \geq -c(1 + |x|^A + |y|^B),$$

(B2), (C1).

Then the boundary value problem (1), (2) possesses at least two solutions.

Proof. As $\alpha \leq \beta$, are well ordered lower and upper solutions, assumptions (A2), (B2), (C1) and Theorem 2 implies the existence of a solution x_1 .

On the other hand, $\alpha_1 \not\leq \beta$, are unordered lower and upper solutions of (1), (2) and also of (14). Moreover, each solution x_2 of (14) is $x_2 \geq \alpha$.

We modify (14) replacing $f_{1\alpha}$ by one sided perturbation

$$f_{1\alpha}^* = \begin{cases} f_{1\alpha}(t, r, y) + M_0, & x > r + 1, \\ f_{1\alpha}(t, r, y) + M_0(x - r), & r < x \leq r + 1, \\ f_{1\alpha}(t, x, y), & x \leq r, \end{cases}$$

where r, M_0 have the same sense as in the proof of Theorem 8.

It is easy to prove by the same method as in the case of Theorem 8 that $x(t) < r$ for each solution $x(t)$ of modified problem. Then there exists a second solution x_2 such that $\alpha < x_2$, $\beta \not\leq x_2 \not\leq \alpha_1$. $\square$

In the case when solution are bounded from above by the strict upper solution β the following symmetric lemma and multiplicity theorem hold.

LEMMA 11. *Let β be a strict upper solution of (1), (2).*

Suppose (C1) holds and sets

$$f_{1\beta}(t, x, y) = \begin{cases} f_1(t, x, y) & \text{for } t \in I, x \leq \beta(t), y \in R, \\ f_1(t, \beta(t), y) & \text{for } t \in I, x > \beta(t), y \in R, \end{cases} \quad (15)$$

and

$$f_{2\beta}(t, x, y) = \begin{cases} f_2(t, x, y) & \text{for } t \in I, x \leq \beta(t), y \in R, \\ x f_2(t, \beta(t), y) & \text{for } t \in I, 0 \geq x < \beta(t), y \in R, \\ 0 & \text{for } t \in I, x > 0 \geq \beta(t), y \in R, \\ f_2(t, \beta(t), y) & \text{for } t \in I, x > \beta(t) > 0, y \in R. \end{cases} \quad (16)$$

Then each solution $x(t)$ of the problem

$$\begin{aligned} (\varphi_p(x'))' &= F_\beta(t, x, \varphi_p(x')), \\ x(b) &= \int_0^b x(s) dg(s) - k\varphi_p(x'(b)), \quad x'(0) = 0 \end{aligned} \quad (17)$$

is a solution of (1), (2).

THEOREM 12. *Let*

(A4) $\alpha \leq \beta$, $\beta_1 \leq \beta$, $\alpha \not\leq \beta_1$, α be a strict lower and β , β_1 a strict upper solutions of (1), (2),

(B4) $\exists 0 < A, B < q - 1, c > 0$ such that $\forall t \in I, x, y \in R$,

$$f_1(t, x, y) \geq -c(1 + |x|^A + |y|^B),$$

(B2), (C1).

Then the boundary value problem (1), (2) possesses at least two solutions.

EXAMPLE 13. We consider the boundary value problem

$$\begin{aligned} (\varphi_p(x'))' - k(t, x)\varphi_m(x) + f(t, x) &= h(t), \\ x(1) &= \int_0^1 x(s) dg(s) - k\varphi_p(x'(1)), \quad x'(0) = 0, \end{aligned} \quad (18)$$

where we assume that $k(t, x) \geq 0$ for $t \in I, x, y \in R$ and the function f is such that

$$\lim_{x \rightarrow -\infty} f(t, x) = \infty, \quad \lim_{x \rightarrow \infty} f(t, x) = -\infty,$$

and

$$\exists x_1, x_2, x_1 < x_2 \quad \text{such that} \quad f(t, x_1) < f(t, x_2) \quad \text{on } I.$$

Then the boundary value problem (18) has a solution for each $h(t)$, at least two solutions for

$$h(t), \quad f(t, x_1) \leq h(t) \leq f(t, x_2) \quad \text{on } I,$$

and at least three solutions for

$$h(t), \quad f(t, x_1) < h(t) < f(t, x_2) \quad \text{on } I.$$

MULTIPLE SOLUTIONS FOR BVP WITH P-LAPLACIAN

REFERENCES

- [1] BIHARI, I.: *A generalization of a lemma of Bellman and its applications to uniqueness problems of differential equations*, Acta Math. Acad. Sci. Hung. **7** (1956), 81–94.
- [2] GAINES, R. E.—MAWHIN J.: *Coincidence Degree, and Nonlinear Differential Equations*, in: Lecture Notes in Math., Vol. 568, Springer, New York, 1977.
- [3] KORMAN, PH.: *Remarks on Nagumo's condition*, Portugal. Math. **55** (1998), 1–9.
- [4] RACHŮNKOVÁ, I.: *On the existence of two solutions of the periodic problem for the ordinary second order differential equation*, Nonlinear Anal. **22** (1994), 1315–1322.
- [5] RACHŮNKOVÁ, I.—STANĚK, S.—TVRDÝ, M.: *Singularities and Laplacians in Boundary Value Problems for Nonlinear Ordinary Differential Equations*, in: Handbook of Differential Equations, Vol. 3, Elsevier, New York, 2006, pp. 607–723.
- [6] RUDOLF, B.: *Method of lower and upper solutions for a generalized boundary value problem*, Arch. Math. (Brno) **36** (2000), 595–602.
- [7] RUDOLF, B.: *An existence and multiplicity result for a periodic boundary value problem*, Math. Bohem. **133** (2008), 41–61.
- [8] RUDOLF, B.: *On a boundary value problem for differential equation with p -Laplacian*, Tatra Mt. Math. Publ. **48** (2011), 189–195.

Received August 8, 2012

Department of Mathematics
FEI STU
Mlynská dolina
SK-812-19 Bratislava
SLOVAKIA
E-mail: boris.rudolf@stuba.sk